

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE USE**



SPREADWET® 1000

Active constituent: 1000 g/L Alkoxylated alcohols

Premium quality, biodegradable wetting and spreading agent for use with insecticides, fungicides and herbicides.

GENERAL INSTRUCTIONS:

Spreadwet 1000 is a premium biodegradable nonionic surfactant for agricultural and horticultural sprays. It contains 100% w/v active ingredients suitable for use with fungicides, insecticides, and herbicides. Spreadwet 1000 reduces the surface tension of spray droplets causing them to spread and adhere to the target in a continuous film. This assists penetration and uptake of the spray onto waxy or hairy plant surfaces, fungous lesions, disease infection sites, and insects. Spreadwet 1000 mixes easily with all types of water and will not injure tender foliage if used as directed.

HERBICIDE SPRAYS:

Spreadwet 1000 can be added to many herbicide sprays to increase their efficiency. Wetter should be added only where directed on individual labels, as use with selective herbicides outside label directions may reduce crop selectivity. For foliar-active herbicides spray to the point where leaves and stems are just wet. Spraying beyond this point will encourage run-off and loss of spray chemicals. When spraying hard-to-wet waxy or hairy-leaved plants, increased rates of Spreadwet 1000 can increase activity. When treating dense, tall vegetation with foliar herbicides, it is important to thoroughly wet leaves and stems for efficient herbicide performance.

DISPOSAL:

Avoid release to the environment or contaminating waterways. Dispose of contents/container at an approved waste disposal plant to comply with local, state and federal regulations.

MIXING:

Spreadwet 1000 mixes easily with water and should be added directly to the spray tank after the addition of other components.

HAZARD CLASSES: H302, H315, H318, H373 **HSR002503**

Hazard & Precautionary Statements

DANGER: Harmful if swallowed. Causes skin irritation. Causes serious eye damage. May cause damage to the organs (kidney/liver) through prolonged or repeated exposure by ingestion, and by inhalation.

PRECAUTIONARY STATEMENTS:

Keep out of reach of children. Read label before use. Do not breathe dust / fume / gas / mist / vapours or spray. Wash contaminated skin thoroughly after handling. Wear protective gloves, protective clothing, eye protection, face protection.

RESPONSE:

If medical advice is needed, have product container or label at hand.

If swallowed: immediately call a poison centre/doctor **0800 POISON (0800 764 766)**.

If on skin: Wash with plenty of soap and water. If skin irritation occurs, get medical advice attention. Take off contaminated clothing and wash before re-use.

If in eyes: Rinse carefully with water for several minutes. Remove contact lenses, if present and easy to do so. Continue rinsing.

FIRST AID:

If swallowed, DO NOT induce vomiting. Rinse mouth with water and give water to drink. For advice contact the **National Poisons Centre 0800 POISON (0800 764 766)** or a doctor immediately.

If skin, hair or eye contact occurs, remove contaminated clothing and flush skin, hair and/or eyes with running water for several minutes. Additional information is contained in the safety data sheet which is available from the Reseller at point of purchase or visit www.agrion.co.nz for alternate contact details.

CONDITIONS OF SALE:

As the use of Spreadwet 1000 is beyond the control of the Manufacturer, to the fullest extent permitted by law, Agrion Crop Solutions Limited disclaims all responsibility or liability for any consequence arising from the use of this product other than in strict compliance with the Directions for Use and other label instructions, and solely for the purpose and in the manner indicated.

® Agrion Crop Solutions and Spreadwet are *the registered trademarks of Agrion Crop Solutions Pty Ltd or its Affiliates.*

IN AN EMERGENCY CALL 0800 154 666



DIRECTIONS FOR USE:

SITUATION	APPLICATION RATE	CRITICAL COMMENTS
Horticultural sprays Insecticides and Fungicides Knapsack	25-75 mL/100L 25 mL/100 L 4 mL/15L	Recommended for addition to all horticultural sprays except where formulation has sufficient wetter.
Protected glasshouse Horticultural crops	10 mL/100L	Rate for sensitive crops in protected glasshouse situation.

Weedkilling sprays High volume power spray Low volume boom spray Knapsack	10-50 mL/100L 40- 100 mL/100L 4 mL/15 L	Refer to individual herbicide labels for specific recommendations.
Glyphosate	100-200 mL/100L	In situations where an organosilicone is not specifically required.
Paraquat and Diquat Boom spray Knapsack	25-100 mL/100L 15 mL/ 15L	To achieve maximum efficiency, use the higher rate of Spreadwet 1000 at low water rates to ensure even coverage of plant foliage and to increase absorption.
Soil residual herbicides e.g. Diuron, Simazine, Atrazine	100 mL/100L	When applying soil active herbicides such as Diuron flowable herbicide to standing weeds, add Spreadwet 1000 to thoroughly wet the foliage. Check specific product labels for individual recommendations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.



Agrion Crop Solutions Limited
119 Carbine Road, Mount Wellington
Auckland 1060, New Zealand
Telephone: +64 884 6358
www.agrion.co.nz